



New Mexico Youth Risk and Resiliency Survey (YRRS)
High School Survey Results
Los Alamos County
Grades 9-12, 2017

New Mexico Department of Health
New Mexico Public Education Department
UNM Prevention Research Center



Suggested citation:

Green D, Peñaloza L, and FitzGerald C. 2018. *New Mexico Youth Risk and Resiliency Survey: High School Survey Results 2017, Los Alamos County*. Epidemiology and Response Division, New Mexico Department of Health; School and Family Support Bureau, New Mexico Public Education Department; and University of New Mexico Prevention Research Center.

This publication was produced by the New Mexico Department of Health (NM DOH), the Public Education Department (NM PED), and the University of New Mexico Prevention Research Center. The NM YRRS receives support from the Centers for Disease Control and Prevention in cooperation with the NM PED through Grant number 1U87PS004195-01. For information about administration and methods used in implementation of the New Mexico Youth Risk and Resiliency Survey (NM YRRS), see the website www.youthrisk.org.

ACKNOWLEDGEMENTS

This report is a product of the NM Youth Risk and Resiliency Survey (YRRS), a project that characterizes risk behaviors and resiliency/protective factors among New Mexico youth. The YRRS is a joint project of the New Mexico Department of Health (NMDOH) and the New Mexico Public Education Department (PED), with support and technical assistance from the University of New Mexico Prevention Research Center (UNM PRC); the Office of Substance Abuse Prevention, Behavioral Health Services Division (OSAP-BHSD); the Albuquerque Area Southwest Tribal Epidemiology Center (AASTEC); and the U.S. Centers for Disease Control and Prevention, Division of Adolescent and School Health (CDC-DASH). Gratitude is extended to the individuals listed below for their contribution toward developing and producing this report. (Individuals who served on the NM-YRRS 2017 Steering Committee are identified by an asterisk (*).)

NEW MEXICO DEPARTMENT OF HEALTH

K. Lynn Gallagher, JD, *Secretary, New Mexico Department of Health*
 Michael Landen, MD, MPH, *State Epidemiologist and Director, Epidemiology and Response Division*
 Toby Rosenblatt, MPA, *Bureau Chief, Injury and Behavioral Epidemiology Bureau*
 Dan Green, MPH, *Survey Epidemiologist, Injury and Behavioral Epidemiology Bureau**
 Ihsan Mahdi, *Morbidity Abuse Epidemiologist, Injury and Behavioral Epidemiology Bureau**
 Laura Tomedi, PhD, MPH, *Alcohol Epidemiologist, Injury and Behavioral Epidemiology Bureau**
 James Padilla, MS, *Tobacco Epidemiologist, Tobacco Use Prevention and Control Program**
 James Farmer, MPH, *Director, Office of School and Adolescent Health**

NEW MEXICO PUBLIC EDUCATION DEPARTMENT

Debbie M. Rael, Deputy Secretary, *School Transformation*
 Denise Koscielniak, Director, *Comprehensive School Supports Division*
 Dean Hopper, Director, *Safe and Healthy Schools Bureau*
 Anne Marlow-Geter, Healthy Schools Coordinator, *Safe and Healthy Schools Bureau**

UNIVERSITY OF NEW MEXICO DIVISION FOR PREVENTION AND POPULATION SCIENCES, HEALTH EVALUATION AND RESEARCH TEAM

Linda J. Peñaloza, PhD, *Associate Research Professor**
 Courtney FitzGerald, MSSW, LMSW, MPH, *Associate Scientist III**
 Robyn Viera, MA, *Scientific Research Manager*
 Laura Gutman, PhD, *Multi-Media Development Specialist*

ALBUQUERQUE AREA SOUTHWEST TRIBAL EPIDEMIOLOGY CENTER

Kevin English, DrPh, MPH, *Director*
 Judith Espinoza, MPH, *Epidemiologist**
 Ophelia Spencer, *Tribal Survey Coordinator*

NEW MEXICO HUMAN SERVICES DEPARTMENT, BEHAVIORAL HEALTH SERVICES DIVISION

Karen Cheman, MPH, *Prevention Staff Manager, Office of Substance Abuse Prevention **

The NM YRRS receives support from the Centers for Disease Control and Prevention in cooperation with the NM PED through Grant number 1U87PS004195-01.

Table of Contents

<u>Topic</u>	<u>Page</u>
Participation in Los Alamos County	7
Risk Behaviors at a Glance	8
Highlights	11
Charts	19
Unintentional Injury	20
Injury: Behaviors Associated with Violence	23
Mental Health	28
Tobacco Use	30
Alcohol Use	34
Drug Use	39
Sexual Behavior	44
Nutrition	49
Physical Activity	51
Body Weight	54
Sexual Identity	56
Gender Identity	57
Other Behaviors and Characteristics	58
Resiliency/Protective Factors	61
Relationship Between Selected Risk Behaviors and Resiliency/Protective Factors	65
Appendix A: Questionnaire with Results	79
Appendix B: About this Report	103
Appendix C: Risk Behavior and Resiliency/Protective Factor Definitions	107

New Mexico Youth Risk and Resiliency Survey
Participation in Los Alamos County
High School (Grades 9-12)

The response rate for Los Alamos County was 85%.

A high response rate produces survey results that are more representative of the student population. A response rate of at least 60% allows generalization of results to the entire student body. A response rate of 70% is excellent and allows a high degree of confidence in results. Response rates below 60% are considered low, and caution should be exercised in interpreting results. Low response rates indicate that the data may represent only students who participated in the survey and not necessarily the entire student body.

Profile of students surveyed

	<u>Number of students who responded</u>	<u>Percent (%)</u>
<u>Total</u>	349	(100%)
<u>Gender</u>		
Girls	194	(55.6%)
Boys	155	(44.4%)

Race/Ethnicity

(Totals may be more than 100% because respondents were allowed to choose multiple race/ethnicities)

American Indian or Alaska Native	26	(8.0%)
Asian	16	(4.9%)
Black or African-American	15	(4.6%)
Hispanic	125	(36.0%)
Native Hawaiian or Pacific Islander	9	(2.8%)
White	286	(88.0%)

Grade Level

9th	99	(28.4%)
10th	109	(31.3%)
11th	83	(23.9%)
12th	57	(16.4%)
Other:	0	(0.0%)

New Mexico Youth Risk and Resiliency Survey

Risk Behaviors at a Glance**Los Alamos County****Grades 9-12**

Indicator	Los Alamos County		New Mexico	
Unintentional Injury	%	(95% CI)	%	(95% CI)
Texted or emailed while driving	37.2	(27.3-48.2)	40.0	(38.2-41.9)
Rarely or never wore a bicycle helmet (of those who rode a bicycle)	44.1	(30.5-58.6)	81.9	(79.0-84.5)
Rarely or never wore a seatbelt	4.9	(2.4-9.5)	6.7	(6.1-7.4)
Violence				
In a physical fight	17.4	(13.2-22.6)	25.9	(24.6-27.2)
In a physical fight on school property	6.4	(4.0-9.9)	9.1	(8.3-9.9)
Carried a weapon on school property	6.3	(2.8-13.5)	5.9	(5.1-6.8)
Skipped school because of safety concerns	6.4	(4.1-9.9)	10.1	(9.2-11.1)
Bullied on school property	31.5	(26.7-36.8)	18.0	(17.2-18.9)
Electronically bullied	22.6	(17.2-29.1)	13.7	(13.0-14.5)
Physical dating violence	11.3	(7.1-17.4)	10.7	(9.8-11.7)
Forced to do sexual things (by anyone)	12.8	(10.0-16.2)	10.4	(9.7-11.1)
Mental Health				
Non-suicidal self-injury	24.5	(20.8-28.7)	20.6	(19.6-21.6)
Felt sadness or hopelessness	33.2	(27.7-39.2)	36.6	(35.5-37.7)
Seriously considered suicide	21.2	(17.3-25.6)	18.3	(17.4-19.2)
Planned suicide	18.3	(15.4-21.6)	16.0	(15.1-16.9)
Attempted suicide	7.6	(5.5-10.4)	10.2	(9.3-11.1)
Injured in a suicide attempt	3.5	(2.0-6.2)	3.2	(2.8-3.6)
Tobacco Use				
Current use of any tobacco product	26.1	(18.5-35.5)	33.8	(32.0-35.7)
Current e-cig use	22.8	(16.7-30.4)	26.4	(25.1-27.8)
Current cigarette smoking	10.2	(6.4-15.9)	11.1	(10.0-12.2)
Current hookah use	6.7	(3.5-12.4)	9.5	(8.6-10.5)
Current cigar use	8.1	(5.5-11.6)	10.0	(9.1-11.1)
Current spit tobacco use	6.3	(3.4-11.4)	7.7	(6.9-8.6)
Alcohol Use				
Current drinking	22.0	(16.4-28.8)	27.5	(10.3-12.3)
Binge drinking	8.7	(5.9-12.7)	11.2	(10.3-12.3)
High intensity binge drinking	3.2	(1.7-5.7)	4.3	(3.7-4.9)
First drink before age 13	17.3	(12.9-22.6)	21.4	(20.3-22.5)
Drinking and driving	6.8	(4.0-11.2)	7.0	(6.3-7.9)

New Mexico Youth Risk and Resiliency Survey

Risk Behaviors at a Glance**Los Alamos County****Grades 9-12**

Indicator	Los Alamos County		New Mexico	
Current Drug Use	%	(95% CI)	%	(95% CI)
Marijuana	16.6	(12.5-21.6)	27.9	(26.0-30.0)
Painkillers to get high	6.1	(3.1-11.5)	6.6	(5.9-7.4)
Synthetic marijuana	2.7	(1.1-6.1)	5.5	(4.8-6.2)
Cocaine	1.6	(0.5-5.6)	4.9	(4.3-5.7)
Inhalants	5.5	(3.7-8.1)	4.9	(4.4-5.6)
Methamphetamine	1.0	(0.3-2.8)	3.0	(2.5-3.5)
Heroin	1.3	(0.3-5.9)	2.8	(2.4-3.3)
Physical Activity, Body Weight, and Nutrition				
Overweight	9.1	(6.8-12.1)	16.0	(15.2-16.9)
Obese	7.5	(5.0-11.2)	15.6	(14.6-16.7)
Overweight or obese	16.7	(13.4-20.5)	31.7	(30.2-33.2)
Daily physical activity	22.6	(17.3-29.0)	28.8	(27.7-29.9)
Daily PE in school	7.8	(5.2-11.7)	24.2	(22.1-26.5)
TV viewing three or more hours daily	14.1	(11.1-17.7)	20.7	(19.7-21.8)
Computer/video use three or more hours daily	34.2	(30.3-38.2)	38.4	(37.1-39.6)
Screen time three or more hours daily	48.7	(43.6-54.0)	55.3	(54.1-56.5)
Ate breakfast daily	38.4	(28.7-49.0)	31.5	(29.9-33.1)
Five servings of fruit or vegetables daily	19.7	(15.5-24.8)	19.9	(19.1-20.8)
Daily soda consumption	15.6	(11.8-20.2)	18.8	(17.5-20.1)
Sexual Behavior				
Sexually Active	21.8	(15.0-30.5)	25.9	(24.0-28.0)
Had sexual intercourse with four or more people	7.2	(3.2-15.4)	9.4	(8.4-10.5)
Had sexual intercourse before age 13	2.5	(1.2-5.1)	3.9	(3.4-4.5)
Used a condom*	56.0	(43.8-67.5)	53.1	(50.8-55.4)
Used effective birth control method*	35.0	(21.9-50.9)	31.5	(29.7-33.4)
Used both a condom and effective birth control*	16.5	(7.3-33.0)	10.1	(9.0-11.3)

* Among sexually active students (those who had sexual intercourse within the past three months)

New Mexico Youth Risk and Resiliency Survey (YRRS)
Highlights
Los Alamos County
Grades 9-12, 2017

Highlights from the New Mexico Youth Risk and Resiliency Survey (YRRS)
 Los Alamos County
 High School (Grades-9-12), 2017

Results from the 2017 High School NM YRRS revealed several encouraging trends in New Mexico. These include declining rates of:

- Alcohol use
- Use of tobacco products such as cigarettes, cigars, and hookahs
- Behaviors associated with violence
- Drug use

At the same time, the prevalence of being overweight or obese has been increasing, and the relatively new phenomenon of using e-cigarettes threatens the overall trend of declining tobacco use among youth.

This report puts the findings for New Mexico alongside YRRS results for Los Alamos County.

Please use these results to help make New Mexico and your own community a safer and healthier place for all of our young people. For more YRRS results, see www.youthrisk.org.


Mental Health

Suicide is one of the leading causes of death among adolescents in New Mexico. After sharp decreases in past-year suicide attempts, the rate has not changed substantially since 2009 (2009 = 9.7%; 2017 = 9.4%). The rate for suicided attempts resulting in injury has also been stable since 2009 (2009 = 3.2%; 2017 = 3.2%), after several years of decreasing rates.

Past Year Mental Health Indicators

NM & Los Alamos County


Grades 9-12, 2017



No statistically significant difference

Tobacco Use


Cigarette smoking among NM high school students has been on the decline since 2003. In 2011, a new question was added to the YRRS about using a hookah, or large water pipe, to smoke tobacco. In 2015, two new questions were added about electronic vapor products (e-cigarettes). In 2017, 26.4% of high school students used e-cigarettes within the last 30 days, more than the percentage who currently any of the other tobacco products measured by the YRRS.

Current Tobacco Use**NM & Los Alamos County****Grades 9-12, 2017**

No statistically significant difference

Alcohol Use

Alcohol use by young people is associated with injuries such as motor vehicle crashes, the leading cause of death among adolescents. Alcohol use at an early age is also associated with an increased risk of chronic liver disease and alcohol dependence later in life. In recent years, rates of current drinking, binge drinking, and drinking and driving have been decreasing among young people nationally and in New Mexico. In 2017 New Mexico high school students had the third highest rate of any state participating in the CDC's Youth Risk Behavior Survey (YRBS) for drinking alcohol before the age of 13.


Alcohol Use**NM & Los Alamos County****Grades 9-12, 2017**

No statistically significant difference

Drug Use

Current use of cocaine, methamphetamines, and pain killers to get high have decreased in recent years. Current marijuana use has been stable since 2005. The rate of current heroin use has been between 2.9% and 3.9% over the past decade, showing little change. New Mexico drug use rates are consistently higher than U.S. rates, except for lifetime use of prescription drugs without a doctor's prescription.

Current Drug Use NM & Los Alamos County Grades 9-12, 2017



**Statistically significant difference*


Sexual Behavior

Early and unsafe sexual activity puts teens at risk of unplanned pregnancy and sexually transmitted infections. Measures of sexual activity among NM high school students are similar to those of other states that participated in the YRBS. Very few sexually active teens used both condoms and reliable birth control.

Sexual Behavior

NM & Los Alamos County


Grades 9-12, 2017



Sexual Behaviors among Sexually Active Students




NM & Los Alamos County

Grades 9-12, 2017




Body Weight, Nutrition, and Physical Activity

Poor nutritional behaviors and a lack of physical activity put young people at risk of becoming overweight or obese. Obese and overweight teens are at risk for many chronic diseases that can cause death or disability. The rates of obesity and overweight increased from 2005-2015, but did not increase from 2015-2017.

Obesity and Overweight**NM & Los Alamos County****Grades 9-12, 2017****Nutrition****NM & Los Alamos County****Grades 9-12, 2017****Physical Activity****NM & Los Alamos County****Grades 9-12, 2017**

Injury: Behaviors Associated with Violence

The rates for fighting and fighting on school property have decreased over the past decade. In 2017, a new question was added to the YRRS that asks about sexual assault (being forced to do things of a sexual nature, such as kissing, touching, or being physically forced to have sexual intercourse).

Behaviors Associated with Violence**NM & Los Alamos County****Grades 9-12, 2017**


Experienced physical dating violence in the past 12 months

Los Alamos County Charts


For definitions of risk behaviors, see Appendix C.

Unintentional Injury


Behaviors Associated with Unintentional Injury by Year, Los Alamos County Grades 9-12, 2005-2017




Unintentional Injury
New Mexico and Los Alamos County
Grades 9-12, 2017




By Sex, Los Alamos County




By Grade, Los Alamos County




Unintentional Injury
New Mexico and Los Alamos County
Grades 9-12, 2017



By Sex, Los Alamos County




By Grade, Los Alamos County




Injury: Behaviors Associated with Violence


Behaviors Associated with Violence by Year, Los Alamos County Grades 9-12, 2003-2017




Injury: Behaviors Associated with Violence
New Mexico and Los Alamos County
Grades 9-12, 2017



By Sex, Los Alamos County




By Grade, Los Alamos County




Injury: Bullying and School Violence


Bullying and School Violence by Year, Los Alamos County Grades 9-12, 2003-2017




Injury: Behaviors Associated with Violence
New Mexico and Los Alamos County
Grades 9-12, 2017




By Sex, Los Alamos County




By Grade, Los Alamos County




Injury: Behaviors Associated with Violence and School Safety
New Mexico and Los Alamos County
Grades 9-12, 2017



By Sex, Los Alamos County




By Grade, Los Alamos County



Mental Health

Mental Health Indicators (past 12 months) by Year, Los Alamos County Grades 9-12, 2003-2017




Mental Health


Mental Health Indicators (past 12 months)

New Mexico and Los Alamos County


Grades 9-12, 2017



By Sex, Los Alamos County




By Grade, Los Alamos County



Tobacco Use


Tobacco Use Indicators by Year, Los Alamos County Grades 9-12, 2005-2017




E-cigarette use was added to the YRRS in 2017.

Hookah use was added to the YRRS in 2011.


Tobacco: Cigarette Use
New Mexico and Los Alamos County
Grades 9-12, 2017




By Sex, Los Alamos County




By Grade, Los Alamos County




Tobacco: Other Use and Exposure
New Mexico and Los Alamos County
Grades 9-12, 2017




By Sex, Los Alamos County




By Grade, Los Alamos County




Tobacco: Other Use and Exposure
New Mexico and Los Alamos County
Grades 9-12, 2017



By Sex, Los Alamos County




By Grade, Los Alamos County




Alcohol Use

Alcohol Use Indicators by Year, Los Alamos County Grades 9-12, 2005-2017




*Starting in 2017, binge drinking referred to 5 or more drinks on a single occasion for boys, and 4 or more drinks on a single occasion for girls, in the past 30 days. Before 2015, binge drinking referred to 5 or more drinks on a single occasion in the past 30 days for both boys and girls.


Alcohol Use
New Mexico and Los Alamos County
Grades 9-12, 2017




By Sex, Los Alamos County




By Grade, Los Alamos County




Behaviors of Current Drinkers: How usually got alcohol
New Mexico and Los Alamos County
Grades 9-12, 2017




By Sex, Los Alamos County




By Grade, Los Alamos County




Behaviors of Current Drinkers: Where usually drank alcohol
New Mexico and Los Alamos County
Grades 9-12, 2017




By Sex, Los Alamos County




By Grade, Los Alamos County




Alcohol Use and Injury
New Mexico and Los Alamos County
Grades 9-12, 2017



By Sex, Los Alamos County





By Grade, Los Alamos County




Drug Use


Drug Use Indicators by Year, Los Alamos County Grades 9-12, 2005-2017




Drug Use: Marijuana
New Mexico and Los Alamos County
Grades 9-12, 2017




By Sex, Los Alamos County




By Grade, Los Alamos County




Current (Past 30 day) Drug Use
New Mexico and Los Alamos County
Grades 9-12, 2017




By Sex, Los Alamos County




By Grade, Los Alamos County




Lifetime Drug Use (Ever used in lifetime)
New Mexico and Los Alamos County
Grades 9-12, 2017




By Sex, Los Alamos County




By Grade, Los Alamos County




Access to Drugs
New Mexico and Los Alamos County
Grades 9-12, 2017



By Sex, Los Alamos County




By Grade, Los Alamos County




Sexual Behavior


Sexual Behavior by Year, Los Alamos County Grades 9-12, 2005-2017




Sexual Behavior
New Mexico and Los Alamos County
Grades 9-12, 2017




By Sex, Los Alamos County




By Grade, Los Alamos County




Sexual Behaviors of Sexually Active Students
New Mexico and Los Alamos County
Grades 9-12, 2017




By Sex, Los Alamos County




By Grade, Los Alamos County




Birth Control Methods Used among of Sexually Active Students
New Mexico and Los Alamos County
Grades 9-12, 2017




By Sex, Los Alamos County




By Grade, Los Alamos County




Birth Control Methods Used among of Sexually Active Students
New Mexico and Los Alamos County
Grades 9-12, 2017



By Sex, Los Alamos County




By Grade, Los Alamos County



Effective birth control method = IUD or implant; shot, patch, or birth control ring; or birth control pills.

Nutrition


Nutrition Indicators by Year, Los Alamos County Grades 9-12, 2005-2017




Nutrition

New Mexico and Los Alamos County


Grades 9-12, 2017



By Sex, Los Alamos County





By Grade, Los Alamos County




Physical Activity


Physical Activity Indicators by Year, Los Alamos County Grades 9-12, 2005-2017




Physical Activity
New Mexico and Los Alamos County
Grades 9-12, 2017




By Sex, Los Alamos County




By Grade, Los Alamos County




Physical Activity
New Mexico and Los Alamos County
Grades 9-12, 2017



By Sex, Los Alamos County




By Grade, Los Alamos County




Body Weight


Body Weight
by Year, Los Alamos County
Grades 9-12, 2005-2017




Body Weight
New Mexico and Los Alamos County
Grades 9-12, 2017




By Sex, Los Alamos County




By Grade, Los Alamos County




Sexual Identity
New Mexico and Los Alamos County
Grades 9-12, 2017




By Sex, Los Alamos County




By Grade, Los Alamos County




Gender Identity
New Mexico and Los Alamos County
Grades 9-12, 2017




By Sex, Los Alamos County




By Grade, Los Alamos County




Other Behaviors and Characteristics
New Mexico and Los Alamos County
Grades 9-12, 2017




By Sex, Los Alamos County




By Grade, Los Alamos County




Other Behaviors and Characteristics
New Mexico and Los Alamos County
Grades 9-12, 2017




By Sex, Los Alamos County




By Grade, Los Alamos County




Other Behaviors and Characteristics
New Mexico and Los Alamos County
Grades 9-12, 2017



By Sex, Los Alamos County




By Grade, Los Alamos County




Resiliency/Protective Factors


Resiliency/Protective Factors
New Mexico and Los Alamos County
Grades 9-12, 2017




Grades 9-12, 2017



**New Mexico and Los Alamos County
Grades 9-12, 2017**



Grades 9-12, 2017




Relationship Between Selected Risk Behaviors and Resiliency/Protective Factors

Physical fighting

by Selected Resiliency Factors

Los Alamos County

Grades 9-12, 2017



How to read this chart:


- Of those who said it was 'A little, pretty much, or very much true' that a parent or other adult was interested in their school work, 15.3% were in a physical fight.
- Of those who said it was 'Not true at all' that a parent or other adult was interested in their school work, 46.8% were in a physical fight.

Skipped school because of safety concerns

by Selected Resiliency Factors

Los Alamos County

Grades 9-12, 2017



How to read this chart:


- Of those who said it was 'A little, pretty much, or very much true' that a parent or other adult was interested in their school work, 5.2% skipped school because of safety concerns.
- Of those who said it was 'Not true at all' that a parent or other adult was interested in their school work, 25.1% skipped school because of safety concerns.

Bullied at school

by Selected Resiliency Factors

Los Alamos County

Grades 9-12 2017



How to read this chart:


- Of those who said it was 'A little, pretty much, or very much true' that a parent or other adult was interested in their school work, 30.6% were bullied at school.
- Of those who said it was 'Not true at all' that a parent or other adult was interested in their school work, 42.7% were bullied at school.

Feelings of sadness or hopelessness

by Selected Resiliency Factors

Los Alamos County

Grades 9-12, 2017



How to read this chart:


- Of those who said it was 'A little, pretty much, or very much true' that a parent or other adult was interested in their school work, 32.0% felt sadness or hopelessness.
- Of those who said it was 'Not true at all' that a parent or other adult was interested in their school work, 53.6% felt sadness or hopelessness.

Suicide attempts

by Selected Resiliency Factors

Los Alamos County

Grades 9-12, 2017



How to read this chart:


- Of those who said it was 'A little, pretty much, or very much true' that a parent or other adult was interested in their school work, 7.5% attempted suicide.
- Of those who said it was 'Not true at all' that a parent or other adult was interested in their school work, 10.1% attempted suicide.

Current cigarette smoking

by Selected Resiliency Factors

Los Alamos County

Grades 9-12, 2017



How to read this chart:


- Of those who said it was 'A little, pretty much, or very much true' that a parent or other adult was interested in their school work, 9.9% were current cigarette smokers.
- Of those who said it was 'Not true at all' that a parent or other adult was interested in their school work, 14.9% were current cigarette smokers.

Binge drinking

by Selected Resiliency Factors

Los Alamos County

Grades 9-12, 2017



How to read this chart:


- Of those who said it was 'A little, pretty much, or very much true' that a parent or other adult was interested in their school work, 8.5% were binge drinkers.
- Of those who said it was 'Not true at all' that a parent or other adult was interested in their school work, 13.5% were binge drinkers.

Drinking and driving

by Selected Resiliency Factors

Los Alamos County

Grades 9-12, 2017



How to read this chart:


- Of those who said it was 'A little, pretty much, or very much true' that a parent or other adult was interested in their school work, 5.6% drove when they had been drinking.
- Of those who said it was 'Not true at all' that a parent or other adult was interested in their school work, 25.9% drove when they had been drinking.

Current painkiller use to get high

by Selected Resiliency Factors

Los Alamos County

Grades 9-12, 2017



How to read this chart:


- Of those who said it was 'A little, pretty much, or very much true' that a parent or other adult was interested in their school work, 4.6% used painkillers to get high.
- Of those who said it was 'Not true at all' that a parent or other adult was interested in their school work, 28.7% used painkillers to get high.

Current cocaine use

by Selected Resiliency Factors

Los Alamos County

Grades 9-12, 2017



How to read this chart:


- Of those who said it was 'A little, pretty much, or very much true' that a parent or other adult was interested in their school work, 1.4% used cocaine.
- Of those who said it was 'Not true at all' that a parent or other adult was interested in their school work, 5.5% used cocaine.

Currently sexually active

by Selected Resiliency Factors

Los Alamos County

Grades 9-12, 2017



How to read this chart:


- Of those who said it was 'A little, pretty much, or very much true' that a parent or other adult was interested in their school work, 21.1% were sexually active.
- Of those who said it was 'Not true at all' that a parent or other adult was interested in their school work, 34.7% were sexually active.

Overweight or obese

by Selected Resiliency Factors

Los Alamos County

Grades 9-12, 2017



How to read this chart:


- Of those who said it was 'A little, pretty much, or very much true' that a parent or other adult was interested in their school work, 16.4% were overweight or obese.
- Of those who said it was 'Not true at all' that a parent or other adult was interested in their school work, 21.3% were overweight or obese.

Unstable housing

by Selected Resiliency Factors

Los Alamos County

Grades 9-12, 2017



How to read this chart:

- Of those who said it was 'A little, pretty much, or very much true' that a parent or other adult was interested in their school work, 3.8% were in unstable housing
- Of those who said it was 'Not true at all' that a parent or other adult was interested in their school work, 5.5% were in unstable housing

Appendix A

Questionnaire with Results

2017 High School Questionnaire
NM YRRS
Los Alamos County

<u>Personal Information</u>	<u>Number of responses</u>	<u>%</u>
1. How old are you?		
A. 12 years old or younger	1	0.3%
B. 13 years old	1	0.3%
C. 14 years old	81	23.2%
D. 15 years old	100	28.7%
E. 16 years old	86	24.6%
F. 17 years old	60	17.2%
G. 18 years old or older	20	5.7%
2. What is your sex?		
A. Female	194	55.6%
B. Male	155	44.4%
3. In what grade are you?		
A. 9th grade	99	28.4%
B. 10th grade	109	31.3%
C. 11th grade	83	23.9%
D. 12th grade	57	16.4%
E. Ungraded or other grade	0	0.0%
4. Are you Hispanic or Latino?		
A. Yes	125	36.0%
B. No	222	64.0%
5. What is your race? (Select one or more responses.)		
NOTE - Because students were allowed to select multiple responses, responses may not total to 100%)		
A. American Indian or Alaska Native	26	8.0%
B. Asian	16	4.9%
C. Black or African American	15	4.6%
D. Native Hawaiian or Other Pacific Islander	9	2.8%
E. White	286	88.0%

	<u>Weighted Mean</u>	
6. How tall are you without your shoes on?		
Mean height in meters -	1.7	
Mean height in feet and inches -	5' 7"	
7. How much do you weigh without your shoes on?		
Mean weight in kilograms -	63.0	
Mean weight in pounds -	138.0	
	<u>Weighted %</u>	<u>95% Confidence Interval</u>
8. Which one of these groups best describes you? (Select only one response.)		
A. American Indian or Alaska Native	3.4	(1.9-5.9)
B. Asian	3.6	(2.2-6.0)
C. Black or African American	1.6	(0.8-3.4)
D. Hispanic or Latino	29.3	(20.0-40.6)
E. Native Hawaiian or Other Pacific Islander	0.0	(.-.)
F. White	62.1	(51.3-71.8)
9. What is the highest level of schooling your mother completed?		
A. Completed grade school or less	1.1	(0.4-3.5)
B. Some high school	3.3	(1.9-5.9)
C. Completed high school	6.9	(4.6-10.3)
D. Some college	16.1	(12.0-21.1)
E. Completed college	35.0	(29.2-41.2)
F. Graduate or professional school	29.5	(22.8-37.2)
G. Not sure	8.1	(6.6-10.0)
10. What is the highest level of schooling your father completed?		
A. Completed grade school or less	1.7	(0.8-3.7)
B. Some high school	2.0	(0.6-6.4)
C. Completed high school	9.5	(6.9-13.0)
D. Some college	8.2	(5.7-11.6)
E. Completed college	25.1	(22.2-28.3)
F. Graduate or professional school	40.8	(33.0-49.2)
G. Not sure	12.6	(9.3-17.0)

	<u>Weighted</u> <u>%</u>	<u>95%</u> <u>Confidence</u> <u>Interval</u>
Personal Safety		
The next 6 questions ask about safety.		
11. When you rode a bicycle during the past 12 months, how often did you wear a helmet?		
A. I did not ride a bicycle during the past 12 months	31.1	(25.1-37.9)
B. Never wore a helmet	22.5	(15.9-30.9)
C. Rarely wore a helmet	7.8	(4.6-13.1)
D. Sometimes wore a helmet	9.1	(6.3-13.1)
E. Most of the time wore a helmet	9.5	(6.6-13.4)
F. Always wore a helmet	19.9	(13.9-27.6)
12. How often do you wear a seat belt when riding in a car driven by someone else?		
A. Never	2.1	(0.9-4.7)
B. Rarely	2.8	(1.3-6.0)
C. Sometimes	6.4	(4.5-9.0)
D. Most of the time	19.2	(16.7-22.0)
E. Always	69.5	(63.5-75.0)
13. During the past 30 days, how many times did you ride in a car or other vehicle driven by someone who had been drinking alcohol?		
A. 0 times	83.4	(78.8-87.2)
B. 1 time	6.9	(5.1-9.4)
C. 2 or 3 times	5.7	(3.6-9.0)
D. 4 or 5 times	0.4	(0.1-1.5)
E. 6 or more times	3.5	(2.1-6.0)
14. During the past 30 days, how many times did you drive a car or other vehicle when you had been drinking alcohol?		
A. I did not drive a car or other vehicle during the past 30 days	35.7	(28.9-43.1)
B. 0 times	59.9	(52.4-67.0)
C. 1 time	1.7	(0.7-3.8)
D. 2 or 3 times	0.6	(0.1-2.4)
E. 4 or 5 times	0.4	(0.0-2.9)
F. 6 or more times	1.7	(0.6-4.9)
15. During the past 30 days, on how many days did you text or e-mail while driving a car or other vehicle?		
A. I did not drive a car or other vehicle during the past 30 days	35.5	(28.7-42.9)
B. 0 days	40.5	(34.0-47.4)
C. 1 or 2 days	8.7	(6.5-11.6)
D. 3 to 5 days	3.1	(1.6-6.0)
E. 6 to 9 days	2.7	(1.0-7.0)
F. 10 to 19 days	2.9	(1.3-6.3)
G. 20 to 29 days	2.6	(1.4-4.9)
H. All 30 days	3.9	(1.4-10.6)

	<u>Weighted</u>	<u>95% Confidence</u>
	<u>%</u>	<u>Interval</u>
16. Is there a gun in your home?		
A. Yes	53.5	(47.1-59.7)
B. No	35.3	(30.6-40.3)
C. Not sure	11.2	(8.8-14.2)

Violence-Related Behaviors**The next 9 questions ask about violence-related behaviors.**

17. During the past 30 days, on how many days did you carry a weapon such as a gun, knife, or club?

A. 0 days	76.2	(72.5-79.5)
B. 1 day	4.3	(2.3-8.1)
C. 2 or 3 days	6.7	(4.7-9.6)
D. 4 or 5 days	2.5	(1.4-4.4)
E. 6 or more days	10.2	(7.1-14.4)

18. During the past 12 months, on how many days did you carry a gun? (Do not count the days when you carried a gun only for hunting or for a sport, such as target shooting.)

A. 0 days	96.7	(92.9-98.5)
B. 1 day	1.1	(0.4-3.1)
C. 2 or 3 days	0.0	(.-.)
D. 4 or 5 days	0.3	(0.0-1.9)
E. 6 or more days	1.9	(0.9-4.1)

19. During the past 30 days, on how many days did you carry a weapon such as a gun, knife, or club on school property?

A. 0 days	93.7	(86.5-97.2)
B. 1 day	0.6	(0.2-2.4)
C. 2 or 3 days	2.0	(0.7-5.3)
D. 4 or 5 days	0.3	(0.0-1.9)
E. 6 or more days	3.4	(1.4-8.1)

20. During the past 30 days, on how many days did you not go to school because you felt you would be unsafe at school or on your way to or from school?

A. 0 days	93.6	(90.1-95.9)
B. 1 day	4.1	(2.8-5.8)
C. 2 or 3 days	0.8	(0.3-2.2)
D. 4 or 5 days	0.2	(0.0-1.3)
E. 6 or more days	1.3	(0.5-3.5)

21. During the past 12 months, how many times were you in a physical fight?

A. 0 times	82.6	(77.4-86.8)
B. 1 time	7.8	(4.9-12.3)
C. 2 or 3 times	5.6	(3.5-8.9)
D. 4 or 5 times	1.1	(0.4-3.2)
E. 6 or 7 times	0.0	(.-.)
F. 8 or 9 times	0.2	(0.0-1.7)
G. 10 or 11 times	0.0	(.-.)
H. 12 or more times	2.7	(1.5-4.9)

	<u>Weighted %</u>	<u>95% Confidence Interval</u>
22. During the past 12 months, how many times were you in a physical fight on school property?		
A. 0 times	93.6	(90.1-96.0)
B. 1 time	3.6	(2.0-6.3)
C. 2 or 3 times	0.6	(0.2-2.6)
D. 4 or 5 times	0.3	(0.0-2.7)
E. 6 or 7 times	0.0	(.-.)
F. 8 or 9 times	0.3	(0.0-2.2)
G. 10 or 11 times	0.0	(.-.)
H. 12 or more times	1.5	(0.7-3.2)
23. Have you ever been physically forced to have sexual intercourse when you did not want to?		
A. Yes	7.5	(4.3-13.0)
B. No	92.5	(87.0-95.7)
24. During the past 12 months, how many times did someone you were dating or going out with physically hurt you on purpose? (Count such things as being hit, slammed into something, or injured with an object or weapon.)		
A. I did not date or go out with anyone during the past 12 months	42.8	(37.2-48.6)
B. 0 times	50.8	(46.2-55.4)
C. 1 time	2.8	(1.3-5.8)
D. 2 or 3 times	2.0	(0.9-4.3)
E. 4 or 5 times	0.4	(0.1-1.6)
F. 6 or more times	1.3	(0.5-3.1)
25. During the past 12 months, how many times did anyone force you to do sexual things that you did not want to do? (Count such things as kissing, touching, or being physically forced to have sexual intercourse.)		
A. I did not date or go out with anyone during the past 12 months		
B. 0 times	87.2	(83.8-90.0)
C. 1 time	4.9	(2.7-8.7)
D. 2 or 3 times	3.6	(2.2-5.9)
E. 4 or 5 times	1.6	(0.8-3.2)
F. 6 or more times	2.7	(1.2-5.6)

Bullying

The next 2 questions ask about bullying. Bullying is when 1 or more students tease, threaten, spread rumors about, hit, shove, or hurt another student over and over again. It is not bullying when 2 students of about the same strength or power argue or fight or tease each other in a friendly way.

26. During the past 12 months, have you ever been bullied on school property?		
A. Yes	31.5	(26.7-36.8)
B. No	68.5	(63.2-73.3)
27. During the past 12 months, have you ever been electronically bullied? (Count being bullied through texting, Instagram, Facebook, or other social media.)		
A. Yes	22.6	(17.2-29.1)
B. No	77.4	(70.9-82.8)

	<u>Weighted</u> <u>%</u>	<u>95%</u> <u>Confidence</u> <u>Interval</u>
Self-Injury		
The next question asks about hurting yourself on purpose.		
28. During the past 12 months, how many times did you do something to purposely hurt yourself without wanting to die, such as cutting or burning yourself on purpose?		
A. 0 times	75.5	(71.3-79.2)
B. 1 time	4.1	(2.8-5.9)
C. 2 or 3 times	8.1	(5.8-11.1)
D. 4 or 5 times	4.6	(3.3-6.5)
E. 6 or more times	7.8	(5.3-11.4)
Sadness and Attempted Suicide		
The next 5 questions ask about sad feelings and attempted suicide. Sometimes people feel so depressed about the future that they may consider attempting suicide, that is, taking some action to end their own lives.		
29. During the past 12 months, did you ever feel so sad or hopeless almost every day for two weeks or more in a row that you stopped doing some usual activities?		
A. Yes	33.2	(27.7-39.2)
B. No	66.8	(60.8-72.3)
30. During the past 12 months, did you ever seriously consider attempting suicide?		
A. Yes	21.2	(17.3-25.6)
B. No	78.8	(74.4-82.7)
31. During the past 12 months, did you make a plan about how you would attempt suicide?		
A. Yes	18.3	(15.4-21.6)
B. No	81.7	(78.4-84.6)
32. During the past 12 months, how many times did you actually attempt suicide?		
A. 0 times	92.4	(89.6-94.5)
B. 1 time	3.8	(2.5-5.7)
C. 2 or 3 times	3.2	(2.0-5.0)
D. 4 or 5 times	0.2	(0.0-1.5)
E. 6 or more times	0.4	(0.1-2.5)
33. If you attempted suicide during the past 12 months, did any attempt result in an injury, poisoning, or overdose that had to be treated by a doctor or nurse?		
A. I did not attempt suicide during the past 12 months	92.4	(89.6-94.5)
B. Yes	3.5	(2.0-6.2)
C. No	4.1	(2.5-6.6)

	<u>Weighted</u> <u>%</u>	<u>95%</u> <u>Confidence</u> <u>Interval</u>
Tobacco Use		
The next 11 questions ask about tobacco use.		
34. Have you ever tried cigarette smoking, even one or two puffs?		
A. Yes	22.2	(15.3-31.2)
B. No	77.8	(68.8-84.7)
35. How old were you when you first tried cigarette smoking, even one or two puffs?		
A. I have never tried cigarette smoking, not even one or two puffs	77.9	(69.0-84.8)
B. 8 years old or younger	1.1	(0.3-3.5)
C. 9 or 10 years old	1.7	(0.6-5.2)
D. 11 or 12 years old	4.8	(2.6-8.7)
E. 13 or 14 years old	7.7	(4.6-12.8)
F. 15 or 16 years old	5.3	(3.4-8.1)
G. 17 years old or older	1.5	(0.6-4.0)
36. During the past 30 days, on how many days did you smoke cigarettes?		
A. 0 days	89.8	(84.1-93.6)
B. 1 or 2 days	5.0	(3.4-7.3)
C. 3 to 5 days	1.9	(0.6-5.5)
D. 6 to 9 days	0.8	(0.2-3.2)
E. 10 to 19 days	0.5	(0.1-1.8)
F. 20 to 29 days	0.2	(0.0-1.3)
G. All 30 days	1.8	(0.5-6.6)
37. During the past 30 days, on the days you smoked, how many cigarettes did you smoke per day?		
A. I did not smoke cigarettes during the past 30 days	89.8	(84.0-93.6)
B. Less than 1 cigarette per day	5.7	(3.2-9.9)
C. 1 cigarette per day	0.8	(0.3-2.1)
D. 2 to 5 cigarettes per day	2.2	(1.1-4.4)
E. 6 to 10 cigarettes per day	0.6	(0.2-2.0)
F. 11 to 20 cigarettes per day	0.3	(0.0-2.2)
G. More than 20 cigarettes per day	0.7	(0.2-2.6)
The next 3 questions ask about electronic vapor products, such as blu, NJOY, Vuse, MarkTen, Logic, Vapin Plus, eGo, and Halo. Electronic vapor products include e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens.		
38. Have you ever used an electronic vapor product?		
A. Yes	37.9	(29.6-46.9)
B. No	62.1	(53.1-70.4)
39. During the past 30 days, on how many days did you use an electronic vapor product?		
A. 0 days	77.2	(69.6-83.3)
B. 1 or 2 days	7.3	(5.1-10.5)
C. 3 to 5 days	4.9	(3.2-7.3)
D. 6 to 9 days	2.4	(1.2-4.5)
E. 10 to 19 days	2.1	(1.1-3.8)
F. 20 to 29 days	2.0	(0.8-5.0)
G. All 30 days	4.1	(2.1-8.1)

	<u>Weighted</u> <u>%</u>	<u>95%</u> <u>Confidence</u> <u>Interval</u>
40. The last time you used an electronic vapor product, what was in the mist or vapor you inhaled?		
A. I have never used an electronic vapor product	65.2	(55.9-73.4)
B. Nicotine	14.0	(10.1-19.1)
C. Marijuana, THC, or hash oil	4.4	(2.5-7.6)
D. Just flavoring	13.2	(8.7-19.6)
E. Something else	0.0	(.-.)
F. Not sure	3.1	(1.5-6.4)
The next 4 questions ask about other tobacco products.		
41. During the past 30 days, on how many days did you use chewing tobacco, snuff, dip, snus, or dissolvable tobacco products, such as Redman, Levi Garrett, Beechnut, Skoal, Skoal Bandits, Copenhagen, Camel Snus, Marlboro Snus, General Snus, Ariva, Stonewall, or Camel Orbs? (Do not count any electronic vapor products.)		
A. 0 days	93.7	(88.6-96.6)
B. 1 or 2 days	2.9	(1.5-5.5)
C. 3 to 5 days	1.2	(0.4-3.8)
D. 6 to 9 days	0.7	(0.3-1.8)
E. 10 to 19 days	0.3	(0.0-2.1)
F. 20 to 29 days	0.0	(.-.)
G. All 30 days	1.2	(0.4-3.2)
42. During the past 30 days, on how many days did you smoke cigars, cigarillos, or little cigars?		
A. 0 days	91.9	(88.4-94.5)
B. 1 or 2 days	3.9	(2.4-6.1)
C. 3 to 5 days	2.2	(1.1-4.6)
D. 6 to 9 days	1.0	(0.3-2.9)
E. 10 to 19 days	0.3	(0.0-2.1)
F. 20 to 29 days	0.0	(.-.)
G. All 30 days	0.7	(0.2-2.6)
43. During the past 30 days, on how many days did you smoke tobacco or flavored tobacco in a hookah, even just a puff?		
A. 0 days	93.3	(87.6-96.5)
B. 1 or 2 days	3.5	(1.7-7.1)
C. 3 to 5 days	1.6	(0.5-4.4)
D. 6 to 9 days	0.7	(0.2-2.2)
E. 10 to 19 days	0.2	(0.0-1.3)
F. 20 to 29 days	0.0	(.-.)
G. All 30 days	0.7	(0.2-2.6)

	<u>Weighted</u> <u>%</u>	<u>95%</u> <u>Confidence</u> <u>Interval</u>
44. During the past 7 days, on how many days were you in the same room with someone who was smoking cigarettes?		
A. 0 days	69.6	(63.2-75.4)
B. 1 day	11.4	(8.9-14.5)
C. 2 days	5.3	(3.3-8.4)
D. 3 days	5.2	(3.4-8.0)
E. 4 days	1.5	(0.8-2.8)
F. 5 days	1.1	(0.5-2.3)
G. 6 days	0.3	(0.1-2.2)
H. 7 days	5.5	(2.6-11.3)

Alcohol

The next 7 questions ask about drinking alcohol. This includes drinking beer, wine, wine coolers, and liquor such as rum, gin, vodka, or whiskey. For these questions, drinking alcohol does not include drinking a few sips of wine for religious purposes.

45. How old were you when you had your first drink of alcohol other than a few sips?		
A. I have never had a drink of alcohol other than a few sips	54.3	(46.1-62.2)
B. 8 years old or younger	4.7	(2.7-8.2)
C. 9 or 10 years old	5.3	(3.1-8.9)
D. 11 or 12 years old	7.2	(5.0-10.3)
E. 13 or 14 years old	12.4	(8.6-17.7)
F. 15 or 16 years old	14.8	(9.9-21.7)
G. 17 years old or older	1.2	(0.4-3.2)
46. During the past 30 days, on how many days did you have at least one drink of alcohol?		
A. 0 days	78.0	(71.2-83.6)
B. 1 or 2 days	9.8	(7.5-12.7)
C. 3 to 5 days	6.6	(4.3-9.9)
D. 6 to 9 days	3.9	(2.3-6.7)
E. 10 to 19 days	0.7	(0.2-2.9)
F. 20 to 29 days	0.0	(.-.)
G. All 30 days	1.1	(0.3-4.4)
47. During the past 30 days, how did you usually get the alcohol you drank?		
A. I did not drink alcohol during the past 30 days	78.9	(71.9-84.5)
B. I bought it in a store such as a liquor store, convenience store, supermarket, discount store, or gas station	0.4	(0.0-2.7)
C. I bought it at a restaurant, bar, or club	0.0	(.-.)
D. I bought it at a public event such as a concert or sporting event	0.0	(.-.)
E. I gave someone else money to buy it for me	3.8	(2.0-7.1)
F. Someone gave it to me	11.2	(7.3-16.7)
G. I took it from a store or family member	2.9	(1.5-5.6)
H. I got it some other way	2.9	(1.5-5.5)

	<u>Weighted</u> <u>%</u>	<u>95%</u> <u>Confidence</u> <u>Interval</u>
48. During the past 30 days, where did you usually drink alcohol? (Select only one response.)		
A. I did not drink alcohol during the past 30 days	76.7	(70.2-82.0)
B. At my home	8.4	(6.5-11.0)
C. At another person's home	11.1	(7.7-15.6)
D. While riding in or driving a car or other vehicle	0.2	(0.0-1.4)
E. At a restaurant, bar, or club	0.3	(0.0-2.2)
F. At a public place such as a park, beach, or parking lot	1.6	(0.6-4.6)
G. At a public event such as a concert or sporting event	0.6	(0.2-2.2)
H. On school property	1.1	(0.4-3.5)
49. During the past 30 days, on how many days did you have at least one drink of alcohol on school property?		
A. 0 days	93.8	(89.0-96.6)
B. 1 or 2 days	3.6	(2.1-6.0)
C. 3 to 5 days	0.9	(0.3-2.3)
D. 6 to 9 days	0.5	(0.1-1.9)
E. 10 to 19 days	0.0	(-.-)
F. 20 to 29 days	0.2	(0.0-1.3)
G. All 30 days	1.0	(0.2-4.1)

The next 2 questions ask about how many drinks of alcohol you have had in a row, that is, within a couple of hours. For the first question, the number of drinks you need to think about is different for female students and male students.

50. During the past 30 days, on how many days did you have 4 or more drinks of alcohol in a row (if you are female) or 5 or more drinks of alcohol in a row (if you are male)?		
A. 0 days	91.3	(87.3-94.1)
B. 1 day	2.2	(1.0-4.8)
C. 2 days	2.9	(1.7-5.1)
D. 3 to 5 days	2.1	(1.1-3.8)
E. 6 to 9 days	0.4	(0.1-1.7)
F. 10 to 19 days	0.3	(0.0-2.2)
G. 20 or more days	0.7	(0.2-2.7)
51. During the past 30 days, what is the largest number of alcoholic drinks you had in a row?		
A. I did not drink alcohol during the past 30 days	79.4	(72.5-85.0)
B. 1 or 2 drinks	8.6	(5.6-13.1)
C. 3 drinks	2.3	(1.6-3.4)
D. 4 drinks	0.2	(0.0-1.4)
E. 5 drinks	1.8	(0.6-5.5)
F. 6 or 7 drinks	2.8	(1.4-5.5)
G. 8 or 9 drinks	1.6	(0.5-5.2)
H. 10 or more drinks	3.2	(1.7-5.7)

	<u>Weighted</u> <u>%</u>	<u>95%</u> <u>Confidence</u> <u>Interval</u>
Marijuana Use		
The next 2 questions ask about marijuana use. Marijuana also is called grass or pot.		
52. How old were you when you tried marijuana for the first time?		
A. I have never tried marijuana	72.9	(65.4-79.4)
B. 8 years old or younger	0.9	(0.3-2.5)
C. 9 or 10 years old	0.8	(0.3-2.2)
D. 11 or 12 years old	3.8	(1.6-8.9)
E. 13 or 14 years old	10.3	(6.7-15.7)
F. 15 or 16 years old	10.9	(8.3-14.0)
G. 17 years old or older	0.3	(0.1-2.0)
53. During the past 30 days, how many times did you use marijuana?		
A. 0 times	83.4	(78.4-87.5)
B. 1 or 2 times	7.1	(5.6-8.9)
C. 3 to 9 times	3.5	(1.9-6.4)
D. 10 to 19 times	0.6	(0.2-2.1)
E. 20 to 39 times	1.1	(0.5-2.8)
F. 40 or more times	4.2	(1.9-9.2)
Other Drugs		
The next 13 questions ask about other drugs.		
54. During your life, how many times have you used any form of cocaine, including powder, crack, or freebase?		
A. 0 times	95.7	(91.1-98.0)
B. 1 or 2 times	1.9	(1.0-3.9)
C. 3 to 9 times	1.0	(0.4-2.4)
D. 10 to 19 times	0.0	(.-.)
E. 20 to 39 times	0.3	(0.0-2.3)
F. 40 or more times	1.0	(0.2-4.1)
55. During the past 30 days, how many times did you use any form of cocaine, including powder, crack, or freebase?		
A. 0 times	98.4	(94.4-99.5)
B. 1 or 2 times	0.3	(0.0-2.1)
C. 3 to 9 times	0.0	(.-.)
D. 10 to 19 times	0.0	(.-.)
E. 20 to 39 times	0.0	(.-.)
F. 40 or more times	1.3	(0.3-5.9)
56. During your life, how many times have you used heroin (also called smack, junk, or China White)?		
A. 0 times	98.4	(94.6-99.5)
B. 1 or 2 times	0.3	(0.0-1.9)
C. 3 to 9 times	0.3	(0.0-2.3)
D. 10 to 19 times	0.0	(.-.)
E. 20 to 39 times	0.3	(0.0-2.3)
F. 40 or more times	0.7	(0.2-2.5)

	<u>Weighted %</u>	<u>95% Confidence Interval</u>
57. During the past 30 days, how many times did you use heroin (also called smack, junk, or China White)?		
A. 0 times	98.7	(94.1-99.7)
B. 1 or 2 times	0.0	(.-.)
C. 3 to 9 times	0.3	(0.0-2.3)
D. 10 to 19 times	0.0	(.-.)
E. 20 to 39 times	0.3	(0.0-2.3)
F. 40 or more times	0.7	(0.2-2.5)
58. During your life, how many times have you used methamphetamines (also called speed, crystal, crank, or ice)?		
A. 0 times	97.4	(94.1-98.9)
B. 1 or 2 times	1.3	(0.6-2.8)
C. 3 to 9 times	0.0	(.-.)
D. 10 to 19 times	0.3	(0.0-2.3)
E. 20 to 39 times	0.3	(0.0-2.3)
F. 40 or more times	0.7	(0.2-2.5)
59. During the past 30 days, how many times did you use methamphetamines (also called speed, crystal, crank, or ice)?		
A. 0 times	99.0	(97.2-99.7)
B. 1 or 2 times	0.3	(0.0-2.1)
C. 3 to 9 times	0.0	(.-.)
D. 10 to 19 times	0.0	(.-.)
E. 20 to 39 times	0.3	(0.0-2.3)
F. 40 or more times	0.3	(0.0-2.5)
60. During your life, how many times have you used ecstasy (also called MDMA)?		
A. 0 times	95.1	(91.4-97.2)
B. 1 or 2 times	2.8	(1.8-4.4)
C. 3 to 9 times	0.2	(0.0-1.5)
D. 10 to 19 times	0.3	(0.0-2.1)
E. 20 to 39 times	0.6	(0.2-2.2)
F. 40 or more times	1.0	(0.2-4.1)
61. During your life, how many times have you taken prescription pain medicine without a doctor's prescription or differently than how a doctor told you to use it? (Count drugs such as codeine, Vicodin, OxyContin, Hydrocodone, and Percocet.)		
A. 0 times	83.9	(78.2-88.4)
B. 1 or 2 times	7.3	(5.0-10.6)
C. 3 to 9 times	4.1	(2.4-7.0)
D. 10 to 19 times	2.3	(1.3-3.9)
E. 20 to 39 times	1.1	(0.4-2.8)
F. 40 or more times	1.4	(0.5-4.0)
62. During your life, how many times have you used a needle to inject any illegal drug into your body?		
A. 0 times	98.8	(96.3-99.6)
B. 1 time	0.6	(0.2-1.9)
C. 2 or more times	0.7	(0.2-2.6)

	<u>Weighted</u> <u>%</u>	<u>95%</u> <u>Confidence</u> <u>Interval</u>
63. During the past 30 days, how many times did you sniff glue, breathe the contents of aerosol spray cans, or inhale any paints or sprays to get high?		
A. 0 times	94.5	(91.9-96.3)
B. 1 or 2 times	4.4	(2.9-6.6)
C. 3 to 9 times	0.2	(0.0-1.3)
D. 10 to 19 times	0.0	(.-.)
E. 20 to 39 times	0.2	(0.0-1.3)
F. 40 or more times	0.7	(0.2-2.6)
64. During the past 30 days, how many times did you use synthetic marijuana (also called K2 or Spice)?		
A. 0 times	97.3	(93.9-98.9)
B. 1 or 2 times	1.2	(0.5-3.0)
C. 3 to 9 times	0.2	(0.0-1.3)
D. 10 to 19 times	0.5	(0.2-1.9)
E. 20 to 39 times	0.3	(0.0-2.5)
F. 40 or more times	0.3	(0.0-2.3)
65. During the past 30 days, how many times did you use a pain killer to get high, like Vicodin, OxyContin (also called Oxy or OC), or Percocet (also called Percs)?		
A. 0 times	93.9	(88.5-96.9)
B. 1 or 2 times	2.8	(1.0-7.3)
C. 3 to 9 times	1.8	(1.1-2.9)
D. 10 to 19 times	0.2	(0.0-1.5)
E. 20 to 39 times	0.3	(0.0-2.1)
F. 40 or more times	1.0	(0.2-4.1)
66. During the past 12 months, has anyone offered, sold, or given you an illegal drug on school property?		
A. Yes	26.2	(20.5-32.9)
B. No	73.8	(67.1-79.5)
Sexual Behavior		
The next 9 questions ask about sexual behavior.		
67. Have you ever had sexual intercourse?	29.4	(21.6-38.6)
A. Yes	70.6	(61.4-78.4)
B. No		
68. How old were you when you had sexual intercourse for the first time?		
A. I have never had sexual intercourse	70.4	(61.1-78.3)
B. 11 years old or younger	1.7	(0.8-3.5)
C. 12 years old	0.9	(0.3-2.5)
D. 13 years old	3.9	(1.7-8.9)
E. 14 years old	5.3	(3.6-7.8)
F. 15 years old	8.5	(5.2-13.7)
G. 16 years old	6.3	(4.0-9.8)
H. 17 years old or older	3.0	(1.1-8.2)

	<u>Weighted</u> <u>%</u>	<u>95%</u> <u>Confidence</u> <u>Interval</u>
69. During your life, with how many people have you had sexual intercourse?		
A. I have never had sexual intercourse	70.9	(61.5-78.9)
B. 1 person	14.4	(11.2-18.4)
C. 2 people	4.0	(1.9-7.9)
D. 3 people	3.5	(2.1-5.7)
E. 4 people	2.4	(0.8-6.9)
F. 5 people	1.3	(0.3-5.3)
G. 6 or more people	3.5	(1.8-6.7)
70. During the past 3 months, with how many people did you have sexual intercourse?		
A. I have never had sexual intercourse	70.9	(61.5-78.9)
B. I have had sexual intercourse, but not during the past 3 months	7.3	(5.6-9.4)
C. 1 person	18.2	(13.1-24.7)
D. 2 people	1.6	(0.6-4.1)
E. 3 people	0.9	(0.3-3.1)
F. 4 people	0.0	(-.-)
G. 5 people	0.0	(-.-)
H. 6 or more people	1.1	(0.4-2.9)
71. Did you drink alcohol or use drugs before you had sexual intercourse the last time?		
A. I have never had sexual intercourse	70.5	(61.2-78.4)
B. Yes	7.9	(5.1-11.9)
C. No	21.6	(15.1-29.8)
72. The last time you had sexual intercourse, did you or your partner use a condom?		
A. I have never had sexual intercourse	71.7	(62.6-79.3)
B. Yes	15.4	(12.6-18.7)
C. No	12.9	(7.6-20.9)
73. The last time you had sexual intercourse, what one method did you or your partner use to prevent pregnancy? (Select only one response.)		
A. I have never had sexual intercourse	71.8	(63.0-79.2)
B. No method was used to prevent pregnancy	3.4	(1.7-6.6)
C. Birth control pills	5.3	(3.8-7.4)
D. Condoms	10.7	(8.3-13.6)
E. An IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon)	3.0	(1.1-8.1)
F. A shot (such as Depo-Provera), patch (such as Ortho Evra), or birth control ring (such as NuvaRing)	0.6	(0.1-2.7)
G. Withdrawal or some other method	4.4	(1.6-11.3)
H. Not sure	0.9	(0.3-2.8)

	<u>Weighted</u> <u>%</u>	<u>95%</u> <u>Confidence</u> <u>Interval</u>
74. During your life, with whom have you had sexual contact?		
A. I have never had sexual contact	53.7	(45.6-61.6)
B. Females	23.9	(19.4-29.1)
C. Males	18.4	(14.8-22.5)
D. Females and males	4.0	(2.4-6.8)
75. Which of the following best describes you?		
A. Heterosexual (straight)	80.8	(73.5-86.4)
B. Gay or lesbian	2.2	(0.9-5.2)
C. Bisexual	10.3	(6.7-15.5)
D. Not sure	6.7	(4.0-11.1)

Food

The next 8 questions ask about food you ate or drank during the past 7 days. Think about all the meals and snacks you had from the time you got up until you went to bed. Be sure to include food you ate at home, at school, at restaurants, or anywhere else.

76. During the past 7 days, how many times did you drink 100% fruit juices such as orange juice, apple juice, or grape juice? (Do not count punch, Kool-Aid, sports drinks, or other fruit-flavored drinks.)

A. I did not drink 100% fruit juice during the past 7 days	41.5	(37.5-45.6)
B. 1 to 3 times during the past 7 days	32.1	(28.6-35.8)
C. 4 to 6 times during the past 7 days	12.1	(9.4-15.4)
D. 1 time per day	5.1	(3.6-7.3)
E. 2 times per day	4.5	(2.6-7.8)
F. 3 times per day	0.6	(0.1-4.6)
G. 4 or more times per day	4.0	(2.3-7.1)

77. During the past 7 days, how many times did you eat fruit? (Do not count fruit juice.)

A. I did not eat fruit during the past 7 days	10.8	(7.7-15.0)
B. 1 to 3 times during the past 7 days	35.6	(29.8-41.7)
C. 4 to 6 times during the past 7 days	18.5	(14.6-23.2)
D. 1 time per day	12.4	(9.7-15.7)
E. 2 times per day	10.6	(7.7-14.4)
F. 3 times per day	6.1	(3.6-10.2)
G. 4 or more times per day	6.0	(3.6-9.9)

78. During the past 7 days, how many times did you eat green salad?

A. I did not eat green salad during the past 7 days	39.3	(34.2-44.7)
B. 1 to 3 times during the past 7 days	37.4	(31.8-43.4)
C. 4 to 6 times during the past 7 days	10.7	(7.7-14.7)
D. 1 time per day	6.6	(3.8-11.2)
E. 2 times per day	2.9	(1.7-5.1)
F. 3 times per day	0.2	(0.0-1.3)
G. 4 or more times per day	2.8	(1.4-5.5)

	<u>Weighted</u> <u>%</u>	<u>95%</u> <u>Confidence</u> <u>Interval</u>
79. During the past 7 days, how many times did you eat potatoes? (Do not count french fries, fried potatoes, or potato chips.)		
A. I did not eat potatoes during the past 7 days	27.9	(24.1-32.0)
B. 1 to 3 times during the past 7 days	52.4	(46.9-57.9)
C. 4 to 6 times during the past 7 days	10.1	(6.7-14.9)
D. 1 time per day	3.9	(2.2-6.8)
E. 2 times per day	2.7	(1.4-5.0)
F. 3 times per day	0.5	(0.1-1.9)
G. 4 or more times per day	2.6	(1.6-4.1)
80. During the past 7 days, how many times did you eat carrots?		
A. I did not eat carrots during the past 7 days	43.7	(36.7-50.9)
B. 1 to 3 times during the past 7 days	37.1	(30.1-44.6)
C. 4 to 6 times during the past 7 days	9.1	(6.2-13.1)
D. 1 time per day	4.5	(3.0-6.7)
E. 2 times per day	2.2	(1.0-5.1)
F. 3 times per day	0.6	(0.2-1.9)
G. 4 or more times per day	2.9	(1.5-5.4)
81. During the past 7 days, how many times did you eat other vegetables? (Do not count green salad, potatoes, or carrots.)		
A. I did not eat other vegetables during the past 7 days	16.1	(11.8-21.6)
B. 1 to 3 times during the past 7 days	36.3	(33.0-39.7)
C. 4 to 6 times during the past 7 days	19.5	(16.4-23.0)
D. 1 time per day	13.4	(9.8-18.1)
E. 2 times per day	7.8	(5.6-10.8)
F. 3 times per day	2.1	(1.1-3.9)
G. 4 or more times per day	4.8	(2.8-8.2)
82. During the past 7 days, how many times did you drink a can, bottle, or glass of soda or pop, such as Coke, Pepsi, or Sprite? (Do not count diet soda or diet pop.)		
A. I did not drink soda or pop during the past 7 days	29.4	(24.6-34.6)
B. 1 to 3 times during the past 7 days	39.1	(33.3-45.1)
C. 4 to 6 times during the past 7 days	16.0	(12.8-20.0)
D. 1 time per day	5.1	(3.3-7.6)
E. 2 times per day	5.6	(4.0-8.0)
F. 3 times per day	1.3	(0.6-2.6)
G. 4 or more times per day	3.6	(2.0-6.3)

	<u>Weighted</u> <u>%</u>	<u>95%</u> <u>Confidence</u> <u>Interval</u>
83. During the past 7 days, on how many days did you eat breakfast?		
A. 0 days	17.3	(13.1-22.5)
B. 1 day	7.6	(4.9-11.6)
C. 2 days	10.2	(8.2-12.6)
D. 3 days	8.7	(5.6-13.2)
E. 4 days	6.3	(3.5-11.0)
F. 5 days	7.2	(5.8-8.9)
G. 6 days	4.4	(3.2-5.9)
H. 7 days	38.4	(28.7-49.0)

Physical Activity**The next 4 questions ask about physical activity.**

84. During the past 7 days, on how many days were you physically active for a total of at least 60 minutes per day? (Add up all the time you spent in any kind of physical activity that increased your heart rate and made you breathe hard some of the time.)

A. 0 days	13.8	(10.2-18.4)
B. 1 day	6.4	(4.1-9.9)
C. 2 days	10.9	(7.8-15.0)
D. 3 days	11.0	(8.2-14.6)
E. 4 days	13.2	(10.2-17.0)
F. 5 days	12.9	(10.7-15.6)
G. 6 days	9.1	(6.8-12.0)
H. 7 days	22.6	(17.3-29.0)
85. On an average school day, how many hours do you watch TV?		
A. I do not watch TV on an average school day	27.8	(22.5-33.7)
B. Less than 1 hour per day	25.8	(21.9-30.1)
C. 1 hour per day	16.7	(12.4-22.1)
D. 2 hours per day	15.7	(12.5-19.6)
E. 3 hours per day	7.2	(5.8-8.9)
F. 4 hours per day	2.8	(1.6-4.7)
G. 5 or more hours per day	4.1	(2.2-7.5)
86. On an average school day, how many hours do you play video or computer games or use a computer for something that is not school work? (Count time spent on things such as Xbox, PlayStation, an iPad or other tablet, a smartphone, texting, YouTube, Instagram, Facebook, or other social media.)		
A. I do not play video or computer games or use a computer for something that is not school work	19.7	(17.1-22.5)
B. Less than 1 hour per day	17.3	(12.9-22.6)
C. 1 hour per day	14.1	(11.9-16.7)
D. 2 hours per day	14.8	(10.6-20.3)
E. 3 hours per day	12.0	(10.1-14.3)
F. 4 hours per day	9.1	(5.8-14.0)
G. 5 or more hours per day	13.0	(9.7-17.3)

	<u>Weighted</u> <u>%</u>	<u>95%</u> <u>Confidence</u> <u>Interval</u>
87. In an average week when you are in school, on how many days do you go to physical education (PE) classes?		
A. 0 days	68.5	(53.1-80.7)
B. 1 day	0.6	(0.2-2.0)
C. 2 days	4.9	(2.2-10.5)
D. 3 days	17.7	(9.8-30.0)
E. 4 days	0.4	(0.1-1.5)
F. 5 days	7.8	(5.2-11.7)

The next question asks about concussions. A concussion is when a blow or jolt to the head causes problems such as headaches, dizziness, being dazed or confused, difficulty remembering or concentrating, vomiting, blurred vision, or being knocked out.

88. During the past 12 months, how many times did you have a concussion from playing a sport or being physically active?

A. 0 times	84.5	(78.5-89.1)
B. 1 time	10.3	(7.5-14.0)
C. 2 times	1.9	(0.8-4.5)
D. 3 times	0.9	(0.2-4.0)
E. 4 or more times	2.3	(1.2-4.5)

Health-Related Topics

The next 10 questions ask about other health-related topics.

89. Have you ever been tested for HIV, the virus that causes AIDS? (Do not count tests done if you donated blood.)

A. Yes	9.6	(7.7-11.9)
B. No	74.3	(70.1-78.2)
C. Not sure	16.1	(12.5-20.4)

90. When was the last time you saw a dentist for a check-up, exam, teeth cleaning, or other dental work?

A. During the past 12 months	86.4	(83.9-88.5)
B. Between 12 and 24 months ago	5.1	(3.6-7.3)
C. More than 24 months ago	3.7	(2.3-5.7)
D. Never	0.6	(0.2-2.4)
E. Not sure	4.2	(2.5-7.0)

91. Has a doctor or nurse ever told you that you have asthma?

A. Yes	26.1	(21.5-31.4)
B. No	70.2	(63.9-75.9)
C. Not sure	3.6	(2.0-6.6)

92. Do you still have asthma?

A. I have never had asthma	55.0	(50.0-60.0)
B. Yes	17.9	(12.9-24.3)
C. No	18.1	(13.5-23.9)
D. Not sure	9.0	(6.2-12.8)

	<u>Weighted</u> <u>%</u>	<u>95%</u> <u>Confidence</u> <u>Interval</u>
93. On an average school night, how many hours of sleep do you get?		
A. 4 or less hours	9.5	(5.1-16.8)
B. 5 hours	11.9	(8.4-16.7)
C. 6 hours	21.7	(19.1-24.5)
D. 7 hours	30.7	(26.3-35.5)
E. 8 hours	20.5	(15.4-26.6)
F. 9 hours	4.8	(3.2-7.2)
G. 10 or more hours	1.0	(0.4-2.6)
94. During the past 12 months, how would you describe your grades in school?		
A. Mostly A's	48.0	(40.8-55.3)
B. Mostly B's	33.1	(28.4-38.1)
C. Mostly C's	11.6	(7.9-16.8)
D. Mostly D's	4.0	(2.6-6.2)
E. Mostly F's	1.6	(0.7-3.8)
F. None of these grades	0.3	(0.0-2.2)
G. Not sure	1.4	(0.5-4.0)
95. Do you have any physical disabilities or long-term health problems? (Long-term means 6-months or more.)		
A. Yes	17.5	(12.8-23.5)
B. No	74.3	(67.9-79.7)
C. Not sure	8.2	(5.2-12.8)
96. During the past 30 days, where did you usually sleep?		
A. In my parent's or guardian's home	96.1	(92.9-97.9)
B. In the home of a friend, family member, or other person because I had to leave my home or my parent or guardian cannot afford housing	1.4	(0.6-3.1)
C. In a shelter or emergency housing	0.3	(0.1-2.2)
D. In a motel or hotel	0.3	(0.0-2.5)
E. In a car, park, campground, or other public place	0.7	(0.2-2.3)
F. I do not have a usual place to sleep	0.3	(0.0-2.2)
G. Somewhere else	0.9	(0.2-3.6)
97. Do you consider yourself transgender, genderqueer, or genderfluid?		
A. No, I am not transgender, genderqueer, or genderfluid	86.7	(83.1-89.6)
B. Yes, I am transgender, genderqueer, or genderfluid	2.3	(1.1-4.9)
C. I am not sure if I am transgender, genderqueer, or genderfluid	2.7	(1.5-4.9)
D. I do not know what this question is asking	8.3	(5.9-11.6)
98. During the past 12 months, how many times have you gambled on a sports team, gambled when playing cards or a dice game, played one of New Mexico's lottery games, gambled on the Internet, or bet on a game of personal skill such as pool or a video game?		
A. 0 times	75.5	(70.0-80.2)
B. 1 or 2 times	12.4	(9.0-17.0)
C. 3 to 9 times	6.3	(3.9-10.1)
D. 10 to 19 times	2.4	(1.1-5.4)
E. 20 to 39 times	0.8	(0.2-3.2)
F. 40 or more times	2.5	(1.1-5.5)

	<u>Weighted</u> <u>%</u>	<u>95%</u> <u>Confidence</u> <u>Interval</u>
You and Your Family		
The next 2 questions ask for some information about you and your family.		
99. How often do you speak a language other than English at home?		
A. Never	68.3	(64.4-72.1)
B. Less than half the time	19.1	(14.5-24.8)
C. About half the time	5.6	(3.5-8.9)
D. More than half the time, but not all of the time	3.6	(1.9-6.6)
E. All of the time	3.4	(2.0-5.7)
100. Were you born in the USA?		
A. Yes	93.5	(91.3-95.2)
B. No	6.5	(4.8-8.7)
Resiliency Factors		
The next 15 questions ask about your family, your school, other adults, your friends, and yourself. How true do you feel the following statements are for you?		
101. In my home, there is a parent or some other adult who is interested in my school work.		
A. Not true at all	6.2	(3.9-9.7)
B. A little true	13.6	(10.7-17.2)
C. Pretty much true	22.1	(18.1-26.7)
D. Very much true	58.1	(53.9-62.1)
102. In my home, there is a parent or some other adult who believes that I will be a success.		
A. Not true at all	4.1	(2.0-8.3)
B. A little true	7.6	(5.7-10.1)
C. Pretty much true	19.1	(16.0-22.6)
D. Very much true	69.1	(63.9-74.0)
103. At my school, there is a teacher or some other adult who listens to me when I have something to say.		
A. Not true at all	6.4	(4.0-10.1)
B. A little true	15.0	(10.5-21.0)
C. Pretty much true	35.8	(32.7-39.0)
D. Very much true	42.7	(37.2-48.4)
104. At my school, there is a teacher or some other adult who believes that I will be a success.		
A. Not true at all	6.1	(3.5-10.5)
B. A little true	10.5	(7.4-14.8)
C. Pretty much true	33.4	(28.6-38.6)
D. Very much true	49.9	(45.7-54.2)
105. Outside of my home and school, there is an adult who really cares about me.		
A. Not true at all	8.9	(5.3-14.6)
B. A little true	10.2	(7.2-14.3)
C. Pretty much true	18.1	(15.1-21.5)
D. Very much true	62.9	(58.3-67.2)

	<u>Weighted</u> <u>%</u>	<u>95%</u> <u>Confidence</u> <u>Interval</u>
106. Outside of my home and school, there is an adult who tells me when I do a good job.		
A. Not true at all	11.6	(8.7-15.4)
B. A little true	17.1	(14.4-20.2)
C. Pretty much true	25.8	(22.1-29.8)
D. Very much true	45.5	(39.4-51.7)
107. I have a friend about my own age who really cares about me.		
A. Not true at all	6.5	(3.6-11.4)
B. A little true	7.9	(5.9-10.5)
C. Pretty much true	18.8	(15.9-22.1)
D. Very much true	66.8	(61.0-72.1)
108. I have a friend about my own age who helps me when I'm having a hard time.		
A. Not true at all	7.6	(4.4-12.9)
B. A little true	11.4	(8.0-16.1)
C. Pretty much true	21.4	(17.8-25.4)
D. Very much true	59.6	(52.2-66.6)
109. When I am not at home, one of my parents/guardians knows where I am and who I am with.		
A. Not true at all	4.4	(2.1-8.9)
B. A little true	12.2	(8.3-17.4)
C. Pretty much true	37.9	(30.2-46.4)
D. Very much true	45.5	(38.5-52.6)
110. In my school, there are clear rules about what students can and cannot do.		
A. Not true at all	4.5	(2.2-9.1)
B. A little true	12.6	(8.4-18.5)
C. Pretty much true	33.6	(27.3-40.5)
D. Very much true	49.3	(42.1-56.5)
111. I plan to go to college or some other school after high school.		
A. Not true at all	3.2	(1.6-6.4)
B. A little true	3.8	(2.3-6.4)
C. Pretty much true	14.7	(10.9-19.5)
D. Very much true	78.3	(70.9-84.2)
112. At school I am involved in sports, clubs, or other extra-curricular activities (such as band, cheerleading, or student council).		
A. Not true at all	20.0	(13.7-28.2)
B. A little true	7.0	(3.8-12.5)
C. Pretty much true	12.3	(8.9-16.8)
D. Very much true	60.7	(50.8-69.8)
113. Outside of my home and school, I am a part of clubs, sports teams, church or temple, or other group activities.		
A. Not true at all	26.2	(22.7-30.0)
B. A little true	12.2	(9.5-15.5)
C. Pretty much true	15.4	(12.0-19.7)
D. Very much true	46.2	(40.5-52.0)

	<u>Weighted</u> <u>%</u>	<u>95%</u> <u>Confidence</u> <u>Interval</u>
114. Outside of my home and school, I am involved in music, art, literature, sports, or a hobby.		
A. Not true at all	17.5	(11.8-25.3)
B. A little true	11.9	(9.4-15.1)
C. Pretty much true	15.8	(13.3-18.8)
D. Very much true	54.7	(46.6-62.5)
115. During the past 30 days, on how many days did you miss classes or school without permission?		
A. 0 days	79.6	(71.9-85.6)
B. 1 or 2 days	13.2	(8.8-19.2)
C. 3 to 5 days	2.9	(1.6-5.3)
D. 6 to 9 days	1.8	(0.7-4.4)
E. 10 or more days	2.5	(1.2-4.9)

This is the end of the survey.

Thank you very much for your help.

Appendix B: About this Report

This report contains results from the 2017 New Mexico High School Youth Risk and Resiliency Survey (YRRS). The YRRS is a statewide survey of public school students in middle schools and high schools throughout New Mexico. Statewide and county level results are available at www.YouthRisk.org and nmhealth.org/go/youth. In addition, school district reports have been prepared for each participating school district and have been distributed to school district offices. School districts are encouraged to reproduce and share these reports with community groups concerned with prevention and health promotion. School district level results will not be released without express permission of the school district superintendent. Forms for requesting school district data are available at www.YouthRisk.org. These documents comprise a useful tool for those working on youth-related issues at the state and local level.

The 2017 YRRS was conducted by selecting two separate samples of student respondents. One sample was selected using a methodology that is highly comparable to that used by other states conducting Youth Risk Behavior Surveys (YRBS), and to the national YRBS methodology. For most reports of YRRS state-level data, that is the sample used to produce results. However, data from that sample cannot be used to represent sub-state areas such as counties or school districts. In order to represent smaller geographical areas, a separate and much larger sample was drawn. This second sample allows generation of estimates for sub-state areas within New Mexico. It is this second sample that was used to generate results for this report. Because of this, rates reported for New Mexico may vary slightly from rates in other reports of state-level YRRS data. The only state-level data reported in this report that were based upon the first sampling methodology are used in the charts below that illustrate how to read error bars. Data in that chart differ slightly from other data in this report.

The YRRS is organized into two major content areas: risk behaviors and protective (resiliency) factors. Risk behaviors include behaviors associated with unintentional injury, violence, mental health, tobacco use, alcohol use, drug use, sexual activity, physical activity, and nutrition. Protective or resiliency factors are measures of the positive and supportive relationships, experiences, activities, resources and values that encourage healthy youth development. Resiliency factors and related traits include caring and supportive relationships in the family, school, community and with peers; boundaries set by the family and school; positive peer support; meaningful participation and constructive use of time in school and outside of school.

Using the Report

The main section of this report consists of a series of charts illustrating results from the 2017 YRRS. Risk behaviors are organized into modules by topic area, with each module starting with one or more line charts demonstrating change in behaviors over the years of the survey. The line charts are followed by a set of bar charts showing a comparison of risk behaviors between the state and the county or school district, a comparison between boys and girls within the county or school district, and a comparison by grade level within the county or school district. For counties or school districts with very few respondents or that haven't participated consistently in the YRRS, some of these charts may be omitted.

The charts about risk behaviors are followed by two sets of charts about resiliency/protective factors. The first of these two sections compares the state rate for each resiliency factor to the county or school district rate. The second section illustrates the relationship between risk behaviors and protective factors. This section consists of a set of bar charts, each showing the prevalence of a particular risk behavior for students with varying levels of selected protective factors.


Understanding the Charts - and a Word about Error Bars

The bar charts in this report present the percentage of students who reported engaging in risk behaviors or who reported having protective factors. The height of the bars conveys the percentage of students reporting each trait, and provides a quick visual comparison between different groups of students reporting that trait. A label for each bar presents the actual percentage of students who reported each trait. For instance, the chart on page 21, Unintentional Injury, shows that 6.0% of students in New Mexico rarely or never wore seatbelts, while 4.9% of students in Los Alamos County rarely or never wore seatbelts.

Because of the high cost and logistical complications involved in conducting a statewide survey, the YRRS was administered to a sample of students, rather than to every single student in each school district. Because YRRS data are from a **sample** of students, and not the entire student body, the results in this report are **estimates**. As with all estimates, there is some uncertainty associated with each of these results. On the charts in this report, this degree of uncertainty is represented by an error bar (confidence interval). The error bar is the thin 'I' shaped line that extends above and below the end of each bar in a chart. One of the major influences on the size of the confidence interval or error bar is the number of participants in the survey. In general, this means that with more students participating in the survey, there will be a higher degree of confidence in the results (i.e., error bars will be smaller).

Interpretation of error bars is illustrated in the charts below. This chart compares the New Mexico prevalence of cigarette smoking and cocaine use to the United States prevalence for 2017. The chart shows that the prevalence of current cigarette smoking was 10.6% in New Mexico and 8.8% in the United States. The error bar for the estimate of current smoking among New Mexico students extends a about 2 percentage points in either direction, from 8.6% to 12.9%. This means that we are confident that the actual percentage of smokers among New Mexico students was between 8.6% to 12.9%. The error bar for the estimate of current smoking in the US extends from 7.2% to 10.7%, and we are confident that the actual percentage of smokers in the United States was between 7.2% and 10.7%.


Ever Used Cocaine and Current Cigarette Smoking
Grades 9-12, NM & US, 2017



How does knowing the upper and lower limits of our error bars help us? While it appears at first glance that New Mexico has a higher prevalence of smoking than the United States (10.6% vs. 8.8%), a closer look at the error bars shows a more complicated picture. The error bars for the New Mexico estimate (8.6%–12.9%) and the US estimate (7.2%–10.7%) overlap. Because of this overlap, we do not have a high degree of confidence that the prevalence of smoking in New Mexico was actually different from the prevalence in the United States. In other words, the difference between the prevalence of smoking in New Mexico and the United States was not statistically significant.

For cocaine use it appears that New Mexico had a higher prevalence than the United States (9.4% vs. 4.8%). The error bar for New Mexico extends from 7.0% to 12.4%, and the error bar for the United States extends from 4.2% to 5.6%. Because the higher bound of the US error bar (5.6%) is lower than the lower bound of the NM error bar (7.0%), these error bars do not overlap. For cocaine use, we are confident in saying that the New Mexico prevalence was higher than the US prevalence. In other words, the difference between the prevalence of cocaine use in New Mexico and the United States was a statistically significant difference.

**Ever Used Cocaine and Current Cigarette Smoking
Grades 9-12, NM & US, 2017**



For some small areas, error bars have been removed from this report. With small populations, the number of students sampled can approach the total student population. In these cases, error bars would have limited meaning because with a high percentage of students surveyed, there is less uncertainty around the estimates than error bars suggest.

Appendix C: Risk Behavior and Resiliency/Protective Factor DefinitionsInjury

Rarely or never wore a bicycle helmet

Answered 'Never' or 'Rarely' to the question, *When you rode a bicycle during the past 12 months, how often did you wear a helmet?*

AMONG those who
reported riding a bicycle

Rarely or never wore a Seatbelt

Answered 'Never' or 'Rarely' to the question, *How often do you wear a seat belt when riding in a car driven by someone else?*

Texted or e-mailed while driving

Answered '1 time' or more to the question, *During the past 30 days, how many times did you text or e-mail while driving a car or other vehicle?*

AMONG those who
drove a car or other vehicle

Has a gun in the home

Answered 'Yes' or more to the question, *Is there a gun in your home?*

Carried a gun

Answered '1 day' or more to the question, *During the past 12 months, on how many days did you carry a gun? (Do not count the days when you carried a gun only for hunting or for a sport, such as target shooting.)*

Violence-Related Behaviors

Carried a weapon

Answered '1 day' or more to the question, *During the past 30 days, on how many days did you carry a weapon such as a gun, knife, or club?*

In a physical fight

Answered '1 time' or more to the question, *During the past 12 months, how many times were you in a physical fight?*

Physical dating violence

Answered '1 time' or more to the question, *During the past 12 months, how many times did someone you were dating or going out with physically hurt you on purpose? (Count such things as being hit, slammed into something, or injured with an object or weapon.)*

Forced to do sexual things (sexual assault)

Answered '1 time' or more to the question, *During the past 12 months, how many times did anyone force you to do sexual things that you did not want to do? (Count such things as kissing, touching, or being physically forced to have sexual intercourse.)*

Ever forced to have sexual intercourse

Answered 'Yes' to the question, *Have you ever been physically forced to have sexual intercourse when you did not want to?*

Electronically bullied

Answered 'Yes' to the question, *During the past 12 months, have you ever been bullied on school property?*

Bullied on school property

Answered 'Yes' to the question, *During the past 12 months, have you ever been electronically bullied? (Include being bullied through e-mail, chat rooms, instant messaging, Web sites, or texting.)*

Carried a weapon on school property

Answered '1 day' or more to the question, *During the past 30 days, on how many days did you carry a weapon such as a gun, knife, or club on school property?*

Skipped school due to safety concerns

Answered '1 day' or more to the question, *During the past 30 days, on how many days did you not go to school because you felt you would be unsafe at school or on your way to or from school?*

In a physical fight on school property

Answered '1 time' or more to the question, *During the past 12 months, how many times were you in a physical fight on school property?*

Mental Health

Non-suicidal self harm

Answered '1 time' or more to the question, *During the past 12 months, how many times did you do something to purposely hurt yourself without wanting to die, such as cutting or burning yourself on purpose?*

Felt sadness or hopelessness

Answered 'Yes' to the question, *During the past 12 months, did you ever feel so sad or hopeless almost every day for two weeks or more in a row that you stopped doing some usual activities?*

Seriously considered suicide

Answered 'Yes' to the question, *During the past 12 months, did you ever seriously consider attempting suicide?*

Made a suicide plan

Answered 'Yes' to the question, *During the past 12 months, did you make a plan about how you would attempt suicide?*

Attempted suicide

Answered '1 time' or more to the question, *During the past 12 months, how many times did you actually attempt suicide?*

Suicide attempt with injury

Answered 'Yes' to the question, *If you attempted suicide during the past 12 months, did any attempt result in an injury, poisoning, or overdose that had to be treated by a doctor or nurse?*

Tobacco Use: Cigarettes

Ever smoked cigarettes

Answered 'Yes' to the question, *Have you ever tried cigarette smoking, even one or two puffs?*

Smoked cigarettes before age 13

Answered '11 or 12 years old' or younger to the question, *How old were you when you first tried cigarette smoking, even one or two puffs?*

Current cigarette smoking (Smoked at least one cigarette in past 30 days)

Answered '1 or 2 days' or more to the question, *During the past 30 days, on how many days did you smoke cigarettes?*

Frequent cigarette smoking (Smoked at least 11 cigarettes per day)

Answered '20 to 29 days' or more to the question, *During the past 30 days, on how many days did you smoke cigarettes?*

Daily cigarette smoking

Answered '30 days' to the question, *How old were you when you first tried cigarette smoking, even one or two puffs?*

Tobacco: Other Use and Exposure

Current cigar use

Answered '1 or 2 days' or more to the question, *During the past 30 days, on how many days did you smoke cigars, cigarillos, or little cigars?*

Current spit tobacco use

Answered '1 or 2 days' or more to the question, *During the past 30 days, on how many days did you use chewing tobacco, snuff, or dip, such as Redman, Levi Garrett, Beechnut, Skoal, Skoal Bandits, or Copenhagen?*

Current e-cigarette use

Answered '1 or 2 days' or more to the question, *During the past 30 days, on how many days did you use an electronic vapor product?*

Ever used e-cigarettes

Answered 'Yes' to the question, *Have you ever used an electronic vapor product?*

Current tobacco use (any of five forms of tobacco)

Any form of tobacco use in the past 30 days (smoked cigarettes or cigars, smoked tobacco with a hookah, used electronic vapor products, or used spit tobacco).

Exposed to second hand cigarette smoke

Answered '1 or 2 days' or more to the question, *During the past 7 days, on how many days were you in the same room with someone who was smoking cigarettes?*

Alcohol

Ever drank alcohol

Gave any response other than *'I have never had a drink of alcohol other than a few sips'* to the question, *How old were you when you had your first drink of alcohol other than a few sips?*

Current drinking

Answered *'1 or 2 days'* or more to the question, *During the past 30 days, on how many days did you have at least one drink of alcohol?*

Binge drinking

Answered *'1 day'* or more to the question, *During the past 30 days, on how many days did you have 4 or more drinks of alcohol in a row (if you are female) or 5 or more drinks of alcohol in a row (if you are male)?*

High intensity binge drinking

Answered *'10 or more drinks'* or more to the question, *During the past 30 days, what is the largest number of alcoholic drinks you had in a row, that is, within a couple of hours?*

First drink before age 13

Answered *'11 or 12 years old or younger'* to the question, *How old were you when you had your first drink of alcohol other than a few sips?*

Rode with drinking driver

Answered *'1 time'* or more to the question, *During the past 30 days, how many times did you ride in a car or other vehicle driven by someone who had been drinking alcohol?*

Drinking and driving

Among those who drove a car, answered *'1 day'* or more to the question, *During the past 30 days, how many times did you drive a car or other vehicle when you had been drinking alcohol?*

Marijuana Use

Ever used marijuana

Gave any response other than *'I have never tried marijuana'* to the question, *How old were you when you tried marijuana for the first time?*

Current marijuana use

Answered *'1 or 2 times'* or more to the question, *During the past 30 days, how many times did you use marijuana?*

Tried marijuana before age 13

Answered *'11 or 12 years old'* or younger to the question, *How old were you when you tried marijuana for the first time?*

Other Drug Use

Current painkiller use to get high

Answered '1 or 2 times' or more to the question, *During the past 30 days, how many times did you use a pain killer to get high, like Vicodin, OxyContin (also called Oxy or OC), or Percocet (also called Percs)?*

Current inhalant use

Answered '1 or 2 times' or more to the question, *During the past 30 days, how many times have you sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high?*

Current cocaine use

Answered '1 or 2 times' or more to the question, *During the past 30 days, how many times did you use any form of cocaine, including powder, crack, or freebase?*

Current ecstasy use

Answered '1 or 2 times' or more to the question, *During the past 30 days, how many times have you used ecstasy?*

Current methamphetamine use

Answered '1 or 2 times' or more to the question, *During the past 30 days, how many times have you used methamphetamines (also called speed, crystal, crank, or ice)?*

Current heroin use

Answered '1 or 2 times' or more to the question, *During the past 30 days, how many times have you used heroin (also called smack, junk, or China White)?*

Prescription drugs without prescription

Answered '1 or 2 times' or more to the question, *During your life, how many times have you taken prescription pain medicine without a doctor's prescription or differently than how a doctor told you to use it? (Count drugs such as codeine, Vicodin, OxyContin, Hydrocodone, and Percocet.)*

Lifetime cocaine use (Ever used in lifetime)

Answered '1 or 2 times' or more to the question, *During your life, how many times did you use any form of cocaine, including powder, crack, or freebase?*

Lifetime ecstasy use (Ever used in lifetime)

Answered '1 or 2 times' or more to the question, *During your life, how many times have you used ecstasy?*

Lifetime methamphetamine use (Ever used in lifetime)

Answered '1 or 2 times' or more to the question, *During your life, how many times have you used methamphetamines (also called speed, crystal, crank, or ice)?*

Lifetime heroin use (Ever used in lifetime)

Answered '1 or 2 times' or more to the question, *During your life, how many times have you used heroin (also called smack, junk, or China White)?*

Injection drug use

Answered '1 or 2 times' or more to the question, *During your life, how many times have you used a needle to inject any illegal drug into your body?*

Offered or sold drugs on school property

Answered 'Yes' to the question, *During the past 12 months, has anyone offered, sold, or given you an illegal drug on school property?*

Sexual Activity

Ever had sexual intercourse

Gave any response other than 'I have never had sexual intercourse' to the question, *How old were you when you had sexual intercourse for the first time?*

Sexually active

Respondent reported having sexual intercourse within the previous 3 months. Answered '1 person' or more to the question, *During the past 3 months, with how many people did you have sexual intercourse?*

Sexual intercourse before age 13

Answered '11 or 12 years old' or younger to the question, *How old were you when you had sexual intercourse for the first time?*

Four+ sexual partners in lifetime

Answered '4 people' or more to the question, *During your life, with how many people have you had sexual intercourse?*
(among all students or among sexually active students, as noted)

Alcohol or drugs before sexual intercourse

Answered 'Yes' to the question, *Did you drink alcohol or use drugs before you had sexual intercourse the last time?*

Used condom

Answered 'Yes' to the question, *The last time you had sexual intercourse, did you or your partner use a condom?*

AMONG those who
reported being "Sexually active" (see above)

Did not use condom

Answered 'No' to the question, *The last time you had sexual intercourse, did you or your partner use a condom?*

AMONG those who
reported being "Sexually active" (see above)

Effective birth control method

Answered 'Birth control pills', 'An IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon)', or 'A shot (such as Depo-Provera), patch (such as Ortho Evra), or birth control ring (such as NuvaRing)' to the question, *The last time you had sexual intercourse, what one method did you or your partner use to prevent pregnancy?*

AMONG those who
reported being "Sexually active" (see above)

Used both a condom and effective birth control

Reported using effective birth control (see above)

AND
used a condom at last sex (see above)

AMONG those who
reported being "Sexually active" (see above)

Sex of sexual contact

Response to the question, *During your life, with whom have you had sexual contact?*

Lesbian, gay, or bisexual

Answered 'Gay or lesbian' or 'Bisexual' to the question, *Which of the following best describes you?*

Nutrition

Five servings of fruits and/or vegetables daily

Reported consuming five or more daily servings of fruit juice, fruit, green salad, potatoes, carrots, or other vegetables over the past 7 days.

Ate three or more vegetables per day

Reported consuming 3 or more daily servings of green salad, potatoes, carrots, or other vegetables.

Daily soda

Answered '1 time per day' or more to the question, *During the past 7 days, how many times did you drink a can, bottle, or glass of soda or pop, such as Coke, Pepsi, or Sprite? (Do not count diet soda or diet pop.)*

Three or more sodas daily

Answered '3 times per day' or more to the question, *During the past 7 days, how many times did you drink a can, bottle, or glass of soda or pop, such as Coke, Pepsi, or Sprite? (Do not count diet soda or diet pop.)*

Ate breakfast daily

Answered '7 days' to the question, *During the past 7 days, on how many days did you eat breakfast?*

Never ate breakfast

Answered '0 days' to the question, *During the past 7 days, on how many days did you eat breakfast?*

Physical Activity

No days of physical activity per week

Answered '0 days' to the question, *During the past 7 days, on how many days were you physically active for a total of at least 60 minutes per day? (Add up all the time you spent in any kind of physical activity that increased your heart rate and made you breathe hard some of the time.)*

Five days of physical activity per week

Answered '5 days' or more to the question, *During the past 7 days, on how many days were you physically active for a total of at least 60 minutes per day? (Add up all the time you spent in any kind of physical activity that increased your heart rate and made you breathe hard some of the time.)*

Daily physical activity

Answered '7 days' to the question, *During the past 7 days, on how many days were you physically active for a total of at least 60 minutes per day? (Add up all the time you spent in any kind of physical activity that increased your heart rate and made you breathe hard some of the time.)*

Watched TV three or more hours per day

Answered '3 hours per day' or more to the question, *On an average school day, how many hours do you watch TV?*

Computer use or video gaming three or more hours per day

Answered '3 hours per day' or more to the question, *On an average school day, how many hours do you play video or computer games or use a computer for something that is not school work? (Count time spent on things such as Xbox, PlayStation, an iPad or other tablet, a smartphone, texting, YouTube, Instagram, Facebook, or other social media.)*

Screen time three or more hours per day

Answered a combined total of '3 hours per day' or more to the two questions above (i.e., TV viewing and or computer/video usage).

No PE in school

Answered '0 days' or more to the question, *In an average week when you are in school, on how many days do you go to physical education (PE) classes?*

Daily PE in school

Answered '5 days' or more to the question, *In an average week when you are in school, on how many days do you go to physical education (PE) classes?*

Concussion

Answered '1 time' or more to the question, *During the past 12 months, how many times did you have a concussion from playing a sport or being physically active?*

Body Weight

Note: Body Mass Index (BMI) was calculated from self reported height and weight.

Overweight

Greater than or equal to the 85th national BMI percentile for age and sex, and less than the 95th national BMI for age and sex, based on historical data.

Obese

Greater than or equal to the 95th national BMI percentile for age and sex, based on historical data.

Overweight or obese

Greater than or equal to the 85th national BMI percentile for age and sex, based on historical data.

Other Behaviors and Characteristics

Physical disability or long-term health problem

Answered 'Yes' the question, *Do you have any physical disabilities or long-term health problems? (Long-term means 6-months or more)*

Ever diagnosed with asthma

Answered 'Yes' the question, *Has a doctor or nurse ever told you that you have asthma?*

Current asthma

Answered 'Yes' the question, *Has a doctor or nurse ever told you that you have asthma?*

AND

Answered 'Yes' the question, *Do you still have asthma?*

Ever had an HIV test

Answered 'Yes' to the question, *Have you ever been tested for HIV, the virus that causes AIDS? (Do not count tests done if you donated blood.)*

Saw a dentist in the last 12 months

Answered 'During the past 12 months' to the question, *When was the last time you saw a dentist for a check-up, exam, teeth cleaning, or other dental work?*

Unstable housing

Gave any response other than 'At home with my parents or guardians' to the question,
During the past 30 days, where did you usually sleep?

Speaks a non-English language

Answered 'More than half the time but not all of the time' OR 'All of the time' to the question, *How often do you speak a language other than English at home?*

Born outside the USA

Answered 'No' to the question, *Were you born in the USA?*

Gambled

Answered '1 or 2 times' or more to the question, *During the past 12 months, how many times have you gambled on a sports team, gambled when playing cards or a dice game, played one of New Mexico's lottery games, gambled on the Internet, or bet on a game of personal skill such as pool or a video game?*

Transgender, genderqueer, or genderfluid

Answered 'Yes, I am transgender, genderqueer, or genderfluid' or more to the question,
Do you consider yourself transgender, genderqueer, or genderfluid?

Cisgender

Answered 'No, I am not transgender, genderqueer, or genderfluid' or more to the question,
Do you consider yourself transgender, genderqueer, or genderfluid?

Resiliency/Protective Factors

These questions are explained in context throughout the report.

